IRSA JOURNAL OF SPORTS STUDIES AND PHYSICAL EDUCATION (IJSSPE)

Volume 1, Issue 1 (2026), pp. 22–25

Open Access Academic Peer-Reviewed Journal

Website: https://ijsspe.oc.com.ar/ijsspe

The Impact of High-Intensity Interval Training (HIIT) on Cardiorespiratory Fitness and Body Composition in Young Football Players

Ali Hassan Al-Musawi[1]*, Mohammed Reza Ahmadi[2]

[1] Department of Physical Education and Sports Sciences, University of Babylon, Iraq

[2] Faculty of Sport Sciences, Bu-Ali Sina University, Hamedan, Iran

**Corresponding Author: ali.almusawi@uobabylon.edu.iq*

Abstract

This study investigates the efficacy of an eight-week High-Intensity Interval Training (HIIT) protocol compared to moderate-intensity continuous training (MICT) on maximum oxygen uptake (VO2max) and body composition metrics among male youth football athletes. Thirty-four participants (N = 34, age 17.2 ± 1.1 years) were randomly assigned to either a HIIT group (4 × 4 min intervals at 90–95% maximum heart rate) or a MICT group (45 minutes continuous running at 65–75% maximum heart rate) performed three times per week alongside routine tactical practice. Pre- and post-intervention evaluations assessed cardiorespiratory endurance via the 20-meter multi-stage fitness test and body fat percentage (BF%) through multi-frequency bioelectrical impedance analysis. The HIIT cohort demonstrated a statistically significant increase in VO2max from 48.6 ± 2.4 to 54.1 ± 2.1 mL/kg/min ($p < 0.01$), outperforming the MICT cohort (48.9 ± 2.2 to 51.0 ± 2.3 mL/kg/min). In addition, body fat percentage decreased markedly in the HIIT cohort (14.8 ± 1.6% to 12.9 ± 1.4%, $p < 0.05$). The findings establish that incorporating short-duration, high-intensity running intervals into in-season training regimens produces superior physiological conditioning and aerobic performance in youth competitive football.

Keywords: *High-Intensity Interval Training, Football, Cardiorespiratory Fitness, Aerobic Endurance, Body Composition.*

1. Introduction

Modern competitive football demands high levels of aerobic capacity and repeated anaerobic sprinting capabilities. Over the course of a standard 90-minute match, professional and youth elite players cover distances ranging between 9 and 12 kilometers at varying physiological intensities. While moderate-intensity continuous training (MICT) has traditionally formed the cornerstone of pre-season conditioning, the dense scheduling of competitive seasons necessitates more time-efficient conditioning strategies.

High-Intensity Interval Training (HIIT), characterized by brief bouts of intense exercise interspersed with recovery periods, has emerged as a potent methodology to induce central and peripheral adaptations. While evidence supports HIIT benefits in adult populations, comparative research assessing its utility against continuous volume training in adolescent developmental football remains an active area of investigation. This study evaluates the specific impact of an eight-week HIIT protocol on cardiorespiratory fitness (VO2max) and body composition parameters in competitive youth football players.

2. Materials and Methods

2.1 Participants and Study Design

A randomized controlled trial design was implemented over eight weeks during the mid-season competitive phase. Thirty-four healthy male outfield football players from regional sports academies participated (N = 34, age: 17.2 ± 1.1 years, body mass: 66.4 ± 5.2 kg, height: 175.6 ± 4.8 cm). Exclusion criteria comprised acute musculoskeletal injuries within the preceding three months or non-compliance exceeding 10% of total intervention sessions. Participants were randomized into two equal groups:

- **HIIT Group (n = 17):** Completed running-based high-intensity interval drills.
- **MICT Group (n = 17):** Completed steady-state moderate aerobic endurance running.

2.2 Training Interventions

Both groups completed three weekly sessions on non-consecutive days following a standardized 15-minute dynamic warm-up:

- **HIIT Protocol:** 4 bouts of 4 minutes running at 90–95% of peak heart rate (HRmax), separated by 3 minutes of active recovery (light jogging at 60–70% HRmax).
- **MICT Protocol:** Continuous running for 45 minutes at a steady intensity corresponding to 65–75% HRmax.

2.3 Outcome Measures

- **Cardiorespiratory Fitness (VO2max):** Estimated using the 20-meter Multi-Stage Shuttle Run Test according to validated standard conversion formulas.
- **Body Composition:** Body mass, body mass index (BMI), and body fat percentage (BF%) were recorded in morning sessions using a calibrated multi-frequency bioelectrical impedance analyzer.

2.4 Statistical Analysis

Data are reported as mean ± standard deviation. Paired sample t-tests assessed within-group pre-to-post changes, and two-way repeated measures analysis of variance (ANOVA) assessed group × time interactions. Statistical significance was designated at $p < 0.05$.

3. Results

All 34 participants completed the eight-week intervention without adverse events.

- The HIIT cohort achieved an 11.3% improvement in VO2max (from 48.6 ± 2.4 to 54.1 ± 2.1 mL/kg/min), significantly surpassing the 4.3% elevation recorded in the MICT group (from 48.9 ± 2.2 to 51.0 ± 2.3 mL/kg/min, $p < 0.01$).
- Body fat percentage decreased significantly in the HIIT group (from 14.8 ± 1.6% down to 12.9 ± 1.4%, $p < 0.01$), whereas the MICT group exhibited minimal change (from 14.6 ± 1.5% to 14.1 ± 1.3%, $p = 0.12$).
- Total body mass remained relatively stable across both cohorts (HIIT: 66.5 ± 5.1 kg vs 65.8 ± 4.9 kg; MICT: 66.3 ± 5.4 kg vs 66.1 ± 5.2 kg), reflecting concurrent muscle preservation.

4. Discussion

The outcomes demonstrate that high-intensity interval conditioning promotes superior aerobic adaptation and lipid profile management relative to moderate continuous running within a constrained training window. The heightened oxygen delivery mechanisms observed following HIIT are predominantly attributed to central cardiovascular adaptations, notably increased stroke volume and left ventricular contractility stimulated by near-maximal training intensities.

From a practical perspective, football coaches face strict time limitations during congested competitive fixtures. HIIT enables equivalent or superior aerobic capacity development in approximately half the training volume required by conventional endurance methods. Incorporating ball-inclusive running drills and interval games may further enrich tactical acumen while achieving targeted cardiovascular conditioning.

5. Conclusion

An eight-week HIIT program yields superior enhancements in cardiorespiratory endurance and positive fat mass reduction compared to traditional continuous aerobic running in adolescent football players. Sports conditioning practitioners should prioritize structured high-intensity intervals to maximize physiological readiness without incurring excessive training volume loads.

References

1. Bangsbo, J., Iaia, F. M., & Krustrup, P. (2008). The Yo-Yo intermittent recovery test: a useful tool for evaluation of physical performance in soccer. *Sports Medicine*, 38(1), 37-51.
2. Buchheit, M., & Laursen, P. B. (2013). High-intensity interval training, solutions to the programming puzzle: Part I: cardiopulmonary emphasis. *Sports Medicine*, 43(5), 313-338.
3. Helgerud, J., Høydal, K., Wang, E., et al. (2007). Aerobic high-intensity intervals improve VO2max more than moderate training. *Medicine & Science in Sports & Exercise*, 39(4), 665-671.
4. Impellizzeri, F. M., Marcora, S. M., et al. (2006). Physiological and performance effects of generic versus specific aerobic training in soccer players. *International Journal of Sports Medicine*, 27(6), 483-492.
5. MacInnis, M. J., & Gibala, M. J. (2017). Physiological adaptations to low-volume high-intensity interval training in human skeletal muscle. *The Journal of Physiology*, 595(9), 2915-2930.